

The tick corrects option are just for idea... Hoped you all make sure by your own self all the answers.... before attempting

**FINAL TERM EXAMINATION**  
**Spring 2010**  
**MTH603- Numerical Analysis (Session - 2)**

**Question No: 1 ( Marks: 1 ) - Please choose one**

Symbol used for forward differences is

- ▶  $\nabla$
- ▶  $\Delta$
- ▶  $\delta$
- ▶  $\mu$

**Question No: 2 ( Marks: 1 ) - Please choose one**

The relationship between central difference operator and the shift operator is given by

- ▶  $\delta = E - E^{-1}$
- ▶  $\delta = E + E^{-1}$
- ▶  $\delta = E^{\frac{1}{2}} + E^{-\frac{1}{2}}$
- ▶  $\delta = E^{\frac{1}{2}} - E^{-\frac{1}{2}}$

**Question No: 3 ( Marks: 1 ) - Please choose one**

Muller's method requires -----starting points

- ▶ 1
- ▶ 2
- ▶ 3
- ▶ 4

Any problem regarding MCS feel free to mail me we try our best level to help you in every steps...

**Question No: 4 ( Marks: 1 ) - Please choose one**

---

If  $S$  is an identity matrix, then

- ▶  $S^{-1} = S$
- ▶  $S^t = S$
- ▶ All are true
- ▶  $S^{-1} = S^t$

**Question No: 5 ( Marks: 1 ) - Please choose one**

---

If we retain  $r+1$  terms in Newton's forward difference formula, we obtain a polynomial of degree ---- agreeing with  $y_x$  at  $x_0, x_1, \dots, x_r$

- ▶  $r+2$
- ▶  $r+1$
- ▶  $r$
- ▶  $r-1$

**Question No: 6 ( Marks: 1 ) - Please choose one**

---

$P$  in Newton's forward difference formula is defined as

- ▶  $p = \left(\frac{x-x_0}{h}\right)$
- ▶  $p = \left(\frac{x+x_0}{h}\right)$
- ▶  $p = \left(\frac{x+x_n}{h}\right)$
- ▶  $p = \left(\frac{x-x_n}{h}\right)$

**Question No: 7 ( Marks: 1 ) - Please choose one**

---

Octal number system has the base -----

- ▶ 2
- ▶ 8

**Any problem regarding MCS feel free to mail me we try our best level to help you in every steps...**

- ▶ 10
- ▶ 16

**Question No: 8 ( Marks: 1 ) - Please choose one**

Newton's divided difference interpolation formula is used when the values of the independent variable are

- ▶ Equally spaced
- ▶ Not equally spaced
- ▶ Constant
- ▶ None of the above

**Question No: 9 ( Marks: 1 ) - Please choose one**

Given the following data

$x$	0	1	2	4
$f(x)$	1	1	2	5

Value of  $f(2,4)$  is

- ▶ 1.5
- ▶ 3
- ▶ 2
- ▶ 1

**Question No: 10 ( Marks: 1 ) - Please choose one**

If  $y(x)$  is approximated by a polynomial  $P_n(x)$  of degree n then the error is given by

- ▶  $\mathcal{E}(x) = y(x) + P_n(x)$
- ▶  $\mathcal{E}(x) = y(x) - P_n(x)$
- ▶  $\mathcal{E}(x) = P_n(x) - y(x)$
- ▶  $\mathcal{E}(x) = y(x) \times P_n(x)$

Any problem regarding MCS feel free to mail me we try our best level to help you in every steps...

**Question No: 11 ( Marks: 1 ) - Please choose one**

Let  $I$  denotes the closed interval spanned by  $x_0, x_1, x_2, x_3, x_4, x_5, x_6, x_7, \bar{x}$ . Then  $F(x)$  vanishes -----times in the interval  $I$ .

- ▶ n-1
- ▶ n+2
- ▶ n
- ▶ n+1

**Question No: 12 ( Marks: 1 ) - Please choose one**

Differential operator in terms of forward difference operator is given by

▶  $D = \frac{1}{h} (\Delta + \frac{\Delta^2}{2!} + \frac{\Delta^3}{3!} + \frac{\Delta^4}{4!} + \frac{\Delta^5}{5!} + \dots)$

▶

$D = \frac{1}{h} (\Delta + \frac{\Delta^2}{2} + \frac{\Delta^3}{3} + \frac{\Delta^4}{4} + \frac{\Delta^5}{5} + \dots)$

▶

$D = \frac{1}{h} (\Delta - \frac{\Delta^2}{2} + \frac{\Delta^3}{3} - \frac{\Delta^4}{4} + \frac{\Delta^5}{5} - \dots)$

▶

$D = \frac{1}{h} (\Delta - \frac{\Delta^2}{2!} + \frac{\Delta^3}{3!} - \frac{\Delta^4}{4!} + \frac{\Delta^5}{5!} - \dots)$

▶

**Question No: 13 ( Marks: 1 ) - Please choose one**

Finding the first derivative of  $f(x)$  at  $x=0.4$  from the following table:

$x$	0.1	0.2	0.3	0.4
$f(x)$	1.10517	1.22140	1.34986	1.49182

Differential operator in terms of -----will be used.

- ▶ Forward difference operator
- ▶ Backward difference operator
- ▶ Central difference operator

**Any problem regarding MCS feel free to mail me we try our best level to help you in every steps...**

- ▶ None of the given choices

**Question No: 14 ( Marks: 1 ) - Please choose one**

For the given table of values

$x$	0.1	0.2	0.3	0.4	0.5	0.6
$f(x)$	0.425	0.475	0.400	0.452	0.525	0.575

$f'(0.1)$ , using two-point equation will be calculated as.....

- ▶ -0.5
- ▶ 0.5
- ▶ 0.75
- ▶ -0.75

**Question No: 15 ( Marks: 1 ) - Please choose one**

In Simpson's 1/3 rule,  $f(x)$  is of the form

- ▶  $ax+b$
- ▶  $ax^2+bx+c$
- ▶  $ax^3+bx^2+cx+d$
- ▶  $ax^4+bx^3+cx^2+dx+e$

**Question No: 16 ( Marks: 1 ) - Please choose one**

$$I = \int_a^b f(x)dx$$

While integrating,  $h$ , width of the interval, is found by the formula--  
---

- ▶  $\frac{b-a}{n}$
- ▶  $\frac{b+a}{n}$

**Any problem regarding MCS feel free to mail me we try our best level to help you in every steps...**

$$\frac{a-b}{n}$$

- ▶
- ▶ None of the given choices

**Question No: 17 ( Marks: 1 ) - Please choose one**

To apply Simpson's 1/3 rule, valid number of intervals are.....

- ▶ 7
- ▶ 8
- ▶ 5
- ▶ 3

**Question No: 18 ( Marks: 1 ) - Please choose one**

For the given table of values

$x$	0.2	0.3	0.4	0.5	0.6	0.7
$f(x)$	0.425	0.475	0.400	0.452	0.525	0.575

$f''(0.2)$ , using three-point equation will be calculated as .....

- ▶ 17.5
- ▶ 12.5
- ▶ 7.5
- ▶ -12.5

**Question No: 19 ( Marks: 1 ) - Please choose one**

To apply Simpson's 1/3 rule, the number of intervals in the following must be

- ▶ 2
- ▶ 3
- ▶ 5
- ▶ 7

**Question No: 20 ( Marks: 1 ) - Please choose one**

To apply Simpson's 3/8 rule, the number of intervals in the following must be

- ▶ 10
- ▶ 11
- ▶ 12
- ▶ 13

Any problem regarding MCS feel free to mail me we try our best level to help you in every steps...

**Question No: 21 ( Marks: 1 ) - Please choose one**

If the root of the given equation lies between a and b, then the first approximation to the root of the equation by bisection method is .....

- ▶  $\frac{(a+b)}{2}$
- ▶  $\frac{(a-b)}{2}$
- ▶  $\frac{(b-a)}{2}$
- ▶ None of the given choices

**Question No: 22 ( Marks: 1 ) - Please choose one**

.....lies in the category of iterative method.

- ▶ Bisection Method
- ▶ Regula Falsi Method
- ▶ Secant Method
- ▶ None of the given choices

**Question No: 23 ( Marks: 1 ) - Please choose one**

For the equation  $x^3 + 3x - 1 = 0$ , the root of the equation lies in the interval.....

- ▶ (1, 3)
- ▶ (1, 2)
- ▶ (0, 1)
- ▶ (1, 2)

**Question No: 24 ( Marks: 1 ) - Please choose one**

Rate of change of any quantity with respect to another can be modeled by

- ▶ An ordinary differential equation
- ▶ A partial differential equation
- ▶ A polynomial equation
- ▶ None of the given choices

**Any problem regarding MCS feel free to mail me we try our best level to help you in every steps...**

**Question No: 25 ( Marks: 1 ) - Please choose one**

---

If  
 $\frac{dy}{dx} = f(x, y)$

Then the integral of this equation is a curve in

- ▶ None of the given choices
- ▶ xt-plane
- ▶ yt-plane
- ▶ xy-plane

**Question No: 26 ( Marks: 1 ) - Please choose one**

---

In solving the differential equation

$$y' = x + y ; y(0.1) = 1.1$$

$h = 0.1$ , By Euler's method  $y(0.2)$  is calculated as

- ▶ 1.44
- ▶ 1.11
- ▶ 1.22
- ▶ 1.33

**Question No: 27 ( Marks: 1 ) - Please choose one**

---

In second order Runge-Kutta method

$k_1$  is given by

- ▶  $k_1 = hf(x_n, y_n)$
- ▶  $k_1 = 2hf(x_n, y_n)$
- ▶  $k_1 = 3hf(x_n, y_n)$
- ▶ None of the given choices

**Question No: 28 ( Marks: 1 ) - Please choose one**

---

In fourth order Runge-Kutta method,  $k_2$  is given by

**Any problem regarding MCS feel free to mail me we try our best level to help you in every steps...**



$$k_2 = hf\left(x_n + \frac{h}{2}, y_n + \frac{k_1}{2}\right)$$



$$k_2 = hf\left(x_n + \frac{h}{3}, y_n + \frac{k_1}{3}\right)$$



$$k_2 = hf\left(x_n - \frac{h}{3}, y_n - \frac{k_1}{3}\right)$$



$$k_2 = hf\left(x_n - \frac{h}{2}, y_n - \frac{k_1}{2}\right)$$



**Question No: 29 ( Marks: 1 ) - Please choose one**

In fourth order Runge-Kutta method,  $k_4$  is given by

$$k_3 = hf(x_n + 2h, y_n + 2k_3)$$



$$k_3 = hf(x_n - h, y_n - k_3)$$



$$k_3 = hf(x_n + h, y_n + k_3)$$



▶ None of the given choices

**Question No: 30 ( Marks: 1 ) - Please choose one**

Adam-Moulton P-C method is derived by employing

- ▶ Newton's backward difference interpolation formula
- ▶ Newton's forward difference interpolation formula
- ▶ Newton's divided difference interpolation formula
- ▶ None of the given choices

**Question No: 31 ( Marks: 2 )**

If  $F(h) = 256.2354$  and  $F\left(\frac{h}{2}\right) = 257.1379$ , then find  $F_1\left(\frac{h}{2}\right)$  using Richardson's extrapolation limit.

**Question No: 32 ( Marks: 2 )**

**Any problem regarding MCS feel free to mail me we try our best level to help you in every steps...**

Evaluate the integral

$$\int_0^{\frac{\pi}{2}} (\cos x + 2) dx$$

Using Simpson's 3/8 rule

$$\frac{\pi}{4}$$

Take h=

**Question No: 33 ( Marks: 2 )**

---

Write a general formula for Modified Euler's method of solving the given differential equation.

**Question No: 34 ( Marks: 3 )**

---

Evaluate the integral

$$\int_0^4 x^2 dx$$

Using Trapezoidal rule

Take h=1

**Question No: 35 ( Marks: 3 )**

---

Evaluate the integral

$$\int_3^5 (\log x + 2) dx$$

Using Simpson's 3/8 rule

Take h=1

**Question No: 36 ( Marks: 3 )**

---

Write a formula for finding the value of  $k_3$  in Fourth-order R-K method.

**Question No: 37 ( Marks: 5 )**

---

Find Newton's forward difference table from the following data.

**Any problem regarding MCS feel free to mail me we try our best level to help you in every steps...**

$x$	0.0	0.1	0.2	0.3	0.4
$f(x)$	1	0.9048	0.8187	0.7408	0.6703

**Question No: 38 ( Marks: 5 )**

---

Evaluate the integral

$$\int_0^3 (x^2 + x) dx$$

Using Simpson's 3/8 rule

Take  $h=1$

**Question No: 39 ( Marks: 5 )**

---

Use Runge-Kutta Method of order four to find the values of  $k_1, k_2, k_3$  and  $k_4$  for the initial value problem

$$y' = \frac{1}{2}(2x^3 + y), y(1) = 2$$

taking  $h = 0.1$

Any problem regarding MCS feel free to mail me we try our best level to help you in every steps...